

# Relative Quadrupole Moments of <sup>192,193</sup>Hg Superdeformed Bands.

*B.C.Busse<sup>1</sup>, P.Fallon<sup>1</sup>, R.Krücken<sup>1</sup>, D.Ackermann<sup>2</sup>, I.Ahmad<sup>2</sup>, S.J.Asztalos<sup>1</sup>,  
D.J.Blumenthal<sup>2</sup>, M.P.Carpenter<sup>2</sup>, R.M.Clark<sup>1</sup>, M.A.Deleplanque<sup>1</sup>, R.M.Diamond<sup>1</sup>,  
S.M.Fischer<sup>2</sup>, F.Hannachi<sup>4</sup>, R.V.F.Janssens<sup>2</sup>, T.L.Khoo<sup>2</sup>, A.Korichi<sup>5</sup>, T.Lauritsen<sup>2</sup>,  
I.Y.Lee<sup>1</sup>, C.J.Lister<sup>2</sup>, A.Lopez-Martens<sup>4</sup>, A.O.Macchiavelli<sup>1</sup>, R.W.MacLeod<sup>1</sup>,  
E.F.Moore<sup>3</sup>, D.Nisius<sup>2</sup>, G.Schmid<sup>1</sup>, D.Seweryniak<sup>2</sup>, F.S.Stephens<sup>1</sup>, K.Vetter<sup>1</sup>*

<sup>1</sup> Lawrence Berkeley Laboratory, Berkeley, California 94720

<sup>2</sup> Argonne National Laboratory, Argonne, Illinois 60439

<sup>3</sup> North Carolina State University, Raleigh, North Carolina 27695

<sup>4</sup> Centre de Spéctrométrie Nucléaire, IN2P3-CNRS bat 104, F-91405, Orsay, France

<sup>5</sup> Institut de Physique Nucléaire, IN2P3-CNRS bat 104, F-91406, Orsay, France

Relative quadrupole moments ( $Q_o$ 's) of yrast and excited SD bands can give insight into the deformation-driving effects of specific nucleon configurations and aid in the interpretation of "identical" bands. A pair of "identical" bands will have transition energies that are equal to, or are at the quarter points or half points of, each other. A clear understanding of this phenomenon has yet to be developed. With this in mind, SD states in <sup>192,193</sup>Hg were populated simultaneously via the <sup>176</sup>Yb(<sup>22</sup>Ne,xn) reaction at a beam energy of 118 MeV. The target consisted of a 500  $\mu\text{g}/\text{cm}^2$  <sup>176</sup>Yb foil evaporated on a 6.8  $\text{mg}/\text{cm}^2$  Au backing. The beam was accelerated by the 88-Inch Cyclotron at Lawrence Berkeley National Laboratory, and  $\gamma$  rays were detected by the Gammasphere array which, for this experiment, had 85 Ge detectors. A total of  $1.9 \times 10^9$  events with a fold  $\geq 5$  were collected. A Doppler Shift Attenuation Method (DSAM) centroid shift analysis was then performed.

Experimental fractional Doppler shifts,  $F(\tau)$ , were extracted and are shown in the figure for the <sup>192</sup>Hg and <sup>193</sup>Hg SD yrast bands. Calculated  $F(\tau)$  curves, that assume a rotational cascade and constant  $Q_o(Q_{sf})$  for the in-band(sidefeeding) states, are shown for comparison. The stopping powers of Ziegler were used. A lineshape analysis on some transitions in the <sup>192,193</sup>Hg SD bands was also performed, leading to results consistent with the  $F(\tau)$  analysis. Comparing the  $F(\tau)$  curves allowed several im-

portant conclusions. The relative  $Q_o$ 's of the yrast SD bands of <sup>192</sup>Hg and <sup>193</sup>Hg are different to a significance  $> 2\sigma$ , with values of  $19.8 \pm 1.2$  eb and  $17.2 \pm .7$  eb respectively. The uncertainty in the sidefeeding time, obtained by  $\chi^2$  minimization, has been included in the  $Q_o$  errors. Nevertheless, radically different sidefeeding in the two SD Hg nuclei could cause the apparent  $Q_o$  difference, although this solution is less likely. Based on current theoretical work, the difference between <sup>192</sup>Hg and <sup>193</sup>Hg is unexpected. The experimental  $Q_o$ 's of all six SD bands in <sup>193</sup>Hg are the same to  $\approx 1\sigma$ , which seems to eliminate a large polarizing effect of the different single particle levels. This similarity in  $Q_o$  is difficult to reconcile, given the difference in the <sup>192</sup>Hg and <sup>193</sup>Hg  $Q_o$ 's. Since <sup>192</sup>Hg and <sup>193</sup>Hg have a SD "identical" band relationship, the data may imply that "identical" transition energies don't always lead to identical deformations. This is the first accurate measurement of  $Q_o$ 's in an odd-mass SD nucleus in the mass 190 region.

